

UNDECAGOLD



95 Horse Block Road, Yaphank NY 11980-9710
Tel: (877) 447-6266 (Toll-Free in US) or (631) 205-9490 Fax: (631) 205-9493
Tech Support: (631) 205-9492 tech@nanoprobes.com
www.nanoprobes.com

PRODUCT INFORMATION

MONOMALEIMIDO UNDECAGOLD LABELING REAGENT

Product Name: MONOMALEIMIDO UNDECAGOLD - 10 nmol
Catalog Number: 2030S (package labeled as 2030A)
Appearance: Orange-yellow powder or solid
Revision: 1.0 (September 2000)

Congratulations on your acquisition of a revolutionary new gold immunoreagent: the MONOMALEIMIDO UNDECAGOLD labeling kit. With this reagent you can label your own primary antibodies (IgG or Fab' fragments), or any other biomolecule containing an accessible sulfhydryl group, with UNDECAGOLD. Because UNDECAGOLD is a discrete molecular compound and not a colloidal gold preparation, conjugates prepared with this reagent have several advantages over colloidal gold conjugates (see below). It is the smallest gold probe commercially available, with a diameter of just 0.8 nm.

CONTENTS

Product Information
Thiol Caution
IgG labeling with UNDECAGOLD
Fab' labeling with UNDECAGOLD
Labeling of other proteins with UNDECAGOLD
General Considerations for Immunostaining with UNDECAGOLD Reagents
Electron Microscopy Immunolabeling with UNDECAGOLD
 1. Cells in Suspension
 2. Thin Sections
Special Considerations for Viewing UNDECAGOLD in the Electron Microscope
Silver enhancement of UNDECAGOLD for Electron Microscopy
Light Microscopy with UNDECAGOLD
Immunoblotting with UNDECAGOLD
References

Warning: For research use only. Not recommended or intended for diagnosis of disease in humans or animals. Do not use internally or externally in humans or animals. Non radioactive and non carcinogenic.

PRODUCT INFORMATION

UNDECAGOLD is the smallest gold label available, prepared using a discrete gold compound rather than a colloid.¹ This kit contains the UNDECAGOLD particle with a single maleimide functionality incorporated into a ligand on the surface of the gold particle; this

has a specific reactivity towards sulfhydryl groups, and may be covalently linked to reduced disulfides in the hinge region of antibodies, as shown on the next page (figure 1). It may also be used to label cysteine residues in proteins, or other biomolecules with an accessible sulfhydryl group. The reagent as supplied has been lyophilized from 0.02 M sodium phosphate at pH 6.5, with 150 mM sodium chloride and 1 mM ethylene diamine tetraacetate; dissolution in 1 ml deionized water will produce a solution of activated UNDECAGOLD in this buffer. UNDECAGOLD conjugates are intended for use in high resolution electron microscopy where the smallest possible probe and the lowest possible interference with the immunoreactivity of the conjugate are desired. They are stable to wide ranges of pH and ionic strength, and are not radioactive or carcinogenic. The labeling reagent should be stored at -20°C.




Figure 1: Schematic showing UNDECAGOLD labeling of Fab' fragment via reaction of a sulfhydryl and a maleimide group.

10 nmol of reagent is supplied: this is sufficient to label up to 0.1 mg of Fab' fragments, or up to 0.3 mg of IgG molecules.

THIOL CAUTION

UNDECAGOLD particles degrade upon exposure to concentrated thiols such as β -mercaptoethanol or dithiothreitol. If such reagents must be used, concentrations should be kept below 1 mM and exposure restricted to 10 minutes or less.

LABELING IgG MOLECULES WITH UNDECAGOLD

IgG molecules contain disulfide bonds which connect the chains in the hinge region. These are reduced with a mild reducing agent, such as mercaptoethylamine hydrochloride (MEA), then reacted with the UNDECAGOLD reagent in buffer solution, either for 1 hour at room temperature or overnight at 4°C. The coupling reaction should be performed at pH 6.5, since at pH values greater than 7 the maleimido group becomes slightly reactive towards primary amines as well as sulfhydryls and may give non-specific labeling. The reduced antibody must be isolated from the MEA before reaction; this may be achieved by gel filtration, using a gel such as Amicon GH-25, which has an exclusion cutoff at molecular weight 3,000. The UNDECAGOLD conjugated product may be isolated either by gel filtration, using a fine gel such as Pharmacia Superose 6 or 12, Superdex 75 or Amicon GCL-90. Alternatively, unbound UNDECAGOLD may be removed by dialysis. The recommended procedure is given below:

1. Dissolve the antibody (0.3 mg IgG) in 0.1 M sodium phosphate buffer, pH 6.0, containing 5 mM EDTA (1 ml), and dissolve MEA (7 mg) in this solution. Incubate at room temperature for 1 hour.
2. Isolate reduced antibody by gel filtration chromatography. Use a gel, such as Amicon GH-25, which has an exclusion cutoff at molecular weight 3,000. As the eluent, use 0.02 M sodium phosphate at pH 6.5, with 150 mM sodium chloride and

1 mM EDTA as preservative. The reduced antibody will be eluted in the void volume as the first sharp peak in the trace. Combine the fractions containing reduced antibody; the total amount of antibody should be calculated from the optical density (usually for IgG, $E_{1\%}$ at 280 nm = 14.5; concentration in mg/ml = $OD_{280\text{ nm}} \times 10/E_{1\%} = OD_{280\text{ nm}} \times 0.69$).

3. Dissolve the UNDECAGOLD reagent in 1 ml deionized water. Sufficient reagent is supplied to label up to 0.3 mg of IgG; if you are using a smaller amount, use a proportionately smaller amount of the UNDECAGOLD reagent (e.g. half the reagent supplied will label 0.15 mg of IgG. Once activated UNDECAGOLD is reconstituted with water it should be used immediately. The maleimide group is hydrolyzed in aqueous solution).
4. Add the activated UNDECAGOLD solution to the reduced antibody. Incubate for 12-18 hours at 4°C.
5. Separate the unbound gold particles from the antibody conjugates using dialysis or gel exclusion chromatography.
 - (a) DIALYSIS: Use a membrane with a molecular weight cutoff of 100,000 and as the exchange buffer use 0.02 M sodium phosphate at pH 7.4 with 150 mM sodium chloride; stir for 4-6 hours in 200 ml of buffer. Repeat twice with fresh buffer. The membrane will retain the UNDECAGOLD labeled antibody, but unbound UNDECAGOLD particles and any other low molecular weight impurities will pass through.
 - (b) GEL FILTRATION: The UNDECAGOLD conjugate may be effectively isolated by HPLC using a gel such as Pharmacia Superose 6 or 12 (which fractionate a wide range of molecular weights) or Amicon GCL-90 (which excludes molecules of mass 30,000 or greater). Concentrate the reaction mixture to a suitably small volume using membrane centrifugation (e.g. Amicon Centricon-30 system). Elute with 0.02 M sodium phosphate at pH 7.4 with 150 mM sodium chloride; the first, faintly colored peak is the conjugate, while the second, darker band is unbound UNDECAGOLD particles. For even higher purity, repeat the this process one time.

The extent of labeling may be determined from the UV/visible spectrum of the conjugate. UNDECAGOLD has an extinction coefficient at 280 nm of 1.7×10^5 , and at 420 nm of 0.47×10^5 ; this means that for IgG labeled 100 % (one UNDECAGOLD particle to one IgG molecule), the ratio of optical densities at 280 nm and 420 nm is close to 8.2.

UNDECAGOLD conjugates should be stored in 0.02 M sodium phosphate buffer with 150 mM sodium chloride; if they are to be stored longer than three days, add 0.1 % bovine serum albumin and 0.05 % sodium azide to prevent bacterial contamination and to prevent the protein from adhering to the surfaces of the storage vessel.

LABELING Fab' FRAGMENTS WITH UNDECAGOLD

If F(ab')₂ fragments are available, they should be used for this purpose. If these are not available, they should be prepared from IgG molecules by pepsin digestion.² It should be noted that IgG molecules from different host animals vary slightly in structure, and therefore will differ in the ease with which they are digested in this way. The procedure given below has been found to be effective in the preparation of F(ab')₂ fragments from IgG developed in goat. Pepsin becomes more active as the pH is lowered; at pH 7 it is inactivated. Some monoclonal antibody subclasses may be unsuitable for this process (for example, IgG3); a comprehensive review has been published which describes the use of pepsin digestion to prepare F(ab')₂ fragments from mouse monoclonal IgG molecules and details which classes may be digested in this manner.

1. Dissolve IgG in 0.1 M sodium citrate buffer at pH 4.5; add a solution of pepsin in 0.5 ml of the same buffer. Use an amount of pepsin equal to 2 % of the mass of IgG. Incubate at 37°C for 20 h using a water bath or incubator.
2. Isolate the F(ab')₂ fragments by HPLC, using a column such as Pharmacia Superose 6 or 12, TSK 3000, or Toyozoda MacMod GF-250 (which have wide molecular weight fractionation ranges) or a superfine exclusion gel such as Amicon GCL-300 (which excludes compounds with molecular weights above 60,000). Elute with 0.02 M sodium phosphate at pH 7.4 with 150 mM sodium chloride and 1 mM EDTA. The F(ab') fragments will elute in the void volume if a GCL-300 column is used, as the first band. Combine the fractions containing F(ab') fragments and calculate the amount using the optical density. Concentrate to 0.5 ml or less using membrane centrifugation.
3. Dilute the F(ab')₂ fragments (0.1 mg) in 0.1 M sodium phosphate buffer, pH 6.0, containing 5 mM EDTA (make up to 1 ml), and dissolve mercaptoethylamine hydrochloride (MEA: 6 mg) in this solution. Incubate at room temperature for 1 hour.
4. Isolate Fab' fragments by gel filtration chromatography. Use a desalting gel, such as Amicon GH-25 which has exclusion cutoff at molecular weight 3,000. As the eluent, use 0.02 M sodium phosphate at pH 6.5, with 150 mM sodium chloride and 1 mM EDTA as preservative. The reduced antibody will be eluted in the void volume as the first sharp peak in the trace. Combine the fractions containing reduced antibody; the total amount of antibody should be calculated from the optical density (usually for Fab', $E_{1\%}$ at 280 nm = 15.3; concentration in mg/ml = $OD_{280\text{ nm}} \times 10/E_{1\%} = OD_{280\text{ nm}} \times 0.65$).

5. Dissolve the UNDECAGOLD reagent in 0.2 ml deionized water. Sufficient reagent is supplied to label up to 0.1 mg of Fab'; if you are using a smaller amount, use a proportionately smaller amount of the UNDECAGOLD reagent (e.g. half the reagent supplied will label 0.05 mg of Fab'. Once activated UNDECAGOLD is reconstituted with water it should be used immediately. The maleimide group is hydrolyzed in aqueous solution).
6. Add the activated UNDECAGOLD solution to the Fab' fragments, and either
 - (a) Incubate at room temperature for 1 hour; or
 - (b) Incubate for 12-18 hours at 4°C.
7. The UNDECAGOLD conjugate may be effectively isolated by HPLC, using a fine gel such as Pharmacia Superose 6 or 12 (which fractionate a wide range of molecular weights) or Amicon GCL-90 (which excludes molecules of mass 30,000 or greater). Concentrate the reaction mixture to a suitably small volume using centrifuge filter (e.g. Amicon Centricon-30) Elute with 0.02 M sodium phosphate at pH 7.4 with 150 mM sodium chloride; the first, faintly colored peak is the conjugate, while the second, darker band is unbound UNDECAGOLD particles. For even higher purity, repeat the process one time.

The extent of labeling may be determined from the UV/visible spectrum of the conjugate. UNDECAGOLD has an extinction coefficient at 280 nm of 1.7×10^5 , and at 420 nm of 0.47×10^5 ; this means that for Fab' labeled 100 % (one UNDECAGOLD particle to one IgG molecule), the ratio of optical densities at 280 nm and 420 nm is close to 5.2.

UNDECAGOLD conjugates should be stored in 0.02 M sodium phosphate buffer with 150 mM sodium chloride at 2 - 8°C; if they are to be stored longer than three days, add 0.1 % bovine serum albumin and 0.05 % sodium azide to prevent bacterial contamination and to prevent the protein from adhering to the surfaces of the storage vessel.

LABELING OTHER PROTEINS WITH UNDECAGOLD

MONOMALEIMIDO UNDECAGOLD may be used to label any protein with an accessible sulfhydryl group, such as a cysteine residue, in the same manner as described above for antibodies.

In some proteins the sulfhydryl functionality is in the form of a disulfide group; this must be reduced with a mild reducing agent, such as MEA, before it can be labeled. If you are unsure of the structure of your protein, and have sufficient quantity available, it is recommended that the suitability of the sulfhydryl for labeling be determined first; some sulfhydryl sites may be buried within the protein structure, and therefore inaccessible to the UNDECAGOLD reagent. The suitability of a particular protein for UNDECAGOLD labeling may be determined using ^{14}C iodoacetic acid before labeling is tried; alternatively, a sensitive colorimetric procedure exists for sulfhydryl determination.³

Sufficient UNDECAGOLD reagent is supplied to label up to 2 nmol of sulfhydryl groups (for example, 0.2 mg of a 100,000 molecular weight compound with one sulfhydryl). Once activated UNDECAGOLD is reconstituted with water it should be used immediately; the maleimide group is hydrolyzed in aqueous solution. The reaction should be performed in the same manner as for antibodies and antibody fragments. For purification and isolation steps alternative buffers may be substituted for those given; however, the labeling reaction itself should be performed using the buffers and conditions specified.

1. If the labeling site is in the form of a disulfide group, it may be reduced to the free sulfhydryl with mercaptoethylamine hydrochloride (MEA). Dissolve the protein in 0.1 M sodium phosphate buffer, pH 6.0, containing 5 mM EDTA (1 ml), and dissolve MEA (8 mg) in this solution. Incubate at room temperature for 1 hour.
2. Isolate reduced protein by gel filtration chromatography. Use a gel, such as Amicon GH-25, which has an exclusion cut-off at molecular weight 3,000. As the eluent, use 0.02 M sodium phosphate at pH 6.5, with 150 mM sodium chloride and 1 mM EDTA as preservative. The reduced antibody will be eluted in the void volume as the first sharp peak in the trace. Combine the fractions containing reduced antibody; the total amount of antibody should be calculated from the optical density.
3. Dissolve the UNDECAGOLD reagent in 1 ml deionized water. Sufficient reagent is supplied to label 1 to 2 nmol of reactive thiols; if you are using a smaller amount, use a proportionately smaller amount of the UNDECAGOLD reagent. Once activated UNDECAGOLD is reconstituted with water it should be used immediately. The maleimide group is hydrolyzed in aqueous solution).
4. Add the activated UNDECAGOLD solution to the reduced protein. Incubate for 12-18 hours at 4°C.
5. Separate the unbound gold particles from the antibody conjugates using dialysis or gel exclusion chromatography.

- (a) DIALYSIS (100,000 OR HIGHER M.W. PROTEINS ONLY): Use a membrane with a molecular weight cutoff of 50,000 or 100,000 and as the exchange buffer use 0.02 M sodium phosphate at pH 7.4 with 150 mM sodium chloride; stir for 4-6 hours in 200 ml of buffer. Repeat twice with fresh buffer. The membrane will retain the UNDECAGOLD labeled antibody, but unbound UNDECAGOLD particles and any other low molecular weight impurities will pass through.
- (b) GEL FILTRATION: The UNDECAGOLD conjugate may be effectively isolated by HPLC using a gel such as Pharmacia Superose 6 or 12 (which fractionate a wide range of molecular weights) or Amicon GCL-90 (which excludes molecules of mass 30,000 or greater). Concentrate the reaction mixture to a suitably small volume using membrane centrifugation (e.g. Amicon Centricon-30 system). Elute with 0.02 M sodium phosphate at pH 7.4 with 150 mM sodium chloride; the first, faintly colored peak is the conjugate, while the second, darker band is unbound UNDECAGOLD particles. For even higher purity, repeat the this process one time.

GENERAL CONSIDERATIONS FOR IMMUNOSTAINING WITH UNDECAGOLD REAGENTS

Basically, normal methodologies may be used successfully with UNDECAGOLD immunoreagents. The concentration of antibody and gold is similar to other commercial preparations of colloidal gold antibodies. Therefore similar dilutions and blocking agents are appropriate.

The major difference will be in the results:

UNDECAGOLD is an extremely uniform 0.8 nm diameter gold particle ($\pm 10\%$).

UNDECAGOLD conjugates contain absolutely no aggregates. This is in sharp contrast to other colloidal gold conjugates that usually are prepared by centrifugation to remove the largest aggregates and frequently contain smaller aggregates.

Close to 1 UNDECAGOLD particle to 1 Fab (or IgG) make this product distinct from the 0.2 - 10 variable stoichiometry of other colloidal gold antibody preparations.

UNDECAGOLD particles do not have affinity to proteins as do other other colloidal golds. This reduces background and false labeling.

USING STAINS WITH UNDECAGOLD

Because the 0.8 nm UNDECAGOLD particles are so small, over staining with OsO_4 , uranyl acetate or lead citrate will obscure direct visualization of individual UNDECAGOLD particles, and therefore these stains should not be used. Only light staining with a low atomic number stain, such as NANOVAN™, a Vanadium based negative stain, should be used.

ELECTRON MICROSCOPY IMMUNOLABELING WITH UNDECAGOLD

If aldehyde-containing reagents have been used for fixation, these must be quenched before labeling. This may be achieved by incubating the specimens for 5 minutes in 50 mM glycine solution in PBS (pH 7.4). Ammonium chloride (50 mM) or sodium borohydride (0.5 - 1 mg/ml) in PBS may be used instead of glycine.

Cells in Suspension

1. Optional fixing of cells: e.g., with glutaraldehyde (0.05 - 1% for 15 minutes) in PBS. Do not use Tris buffer since this contains an amine. After fixation, centrifuge cells (e.g. 1 ml at 10^7 cells/ml) at 300 X g, 5 minutes; discard supernatant; resuspend in 1 ml buffer. Repeat this washing (centrifugation and resuspension) 2 times.
2. Incubate cells with 0.02 M glycine in PBS (5 mins). Centrifuge, then resuspend cells in PBS-BSA buffer (specified below) for 5 minutes.
4. Wash cells using PBS-BSA as described in step 1 (2 X 5 mins). Resuspend in 1 ml Buffer 1.
3. Place 50 - 200 μl of cells into Eppendorf tube. Dilute UNDECAGOLD conjugate ~ 50 times in PBS-BSA buffer and add 30 μl to cells; incubate for 30 minutes with occasional shaking (do not create bubbles which will denature proteins).
5. Wash cells in PBS-BSA as described in step 1 (2 X 5 mins).

6. Fix cells and antibodies using a final concentration of 1% glutaraldehyde in PBS for 15 minutes. Then remove fixative by washing with buffer 1 (3 X 5 mins).

PBS-BSA Buffer:

20 mM phosphate
 150 mM NaCl
 pH 7.4
 0.5% BSA
 0.1% gelatin (high purity)

Optional, may reduce background:

0.5 M NaCl
 0.05% Tween 20

PBS Buffer:

20 mM phosphate
 150 mM NaCl
 pH 7.4

Negative staining may be used for electron microscopy of small structures or single molecules which are not embedded. Negative stain must be applied after the silver enhancement. NANOVAN™ negative stain is specially formulated for use with UNDECAGOLD reagents; it is based on vanadium, which gives a lighter stain than uranium, lead or tungsten-based negative stains and allows easier visualization of UNDECAGOLD particles with little or no silver enhancement.

Thin Sections

Labeling with UNDECAGOLD may be performed before or after embedding.⁴ Labeling before embedding and sectioning (the pre-embedding method) is used for the study of surface antigens, particularly small organisms such as viruses budding from host cells. It gives good preservation of cellular structure, and subsequent staining usually produces high contrast for study of the cellular details. Labeling after embedding and sectioning (the post-embedding method) allows the antibody access to the interior of the cells, and is used to label both exterior and interior features. The procedures for both methods are described below.

Thin sections mounted on grids are floated on drops of solutions on parafilm or in well plates. Hydrophobic resins usually require pre-etching.

PROCEDURE FOR PRE-EMBEDDING METHOD:⁴

1. Float on a drop of water for 5 - 10 minutes.
2. Incubate cells with 1 % bovine serum albumin in PBS buffer at pH 7.4 for 5 minutes; this blocks any non-specific protein binding sites and minimizes non-specific antibody binding.
3. Rinse with PBS-BSA (1 min).
4. Incubate with UNDECAGOLD conjugate diluted 1/40 - 1/200 in PBS-BSA with 1 % normal serum from the same species as the UNDECAGOLD labeled antibody, for 10 minutes to 1 hour at room temperature.
5. Rinse with PBS-BSA (3 X 1 min), then PBS (3 X 1 min).
6. Postfix with 1 % glutaraldehyde in PBS (10 mins).
7. Rinse in deionized water (2 X 5 min).
8. Dehydrate and embed according to usual procedure. Use of a low-temperature resin (eg. Lowicryl) is recommended.
9. Stain (uranyl acetate, lead citrate or other positive staining reagent) as usual before examination.

Silver enhancement may be performed before or after embedding (see below); it should be completed before postfixing or staining with osmium tetroxide, uranyl acetate or similar reagents is carried out.

PROCEDURE FOR POST-EMBEDDING METHOD:⁴

1. Prepare sections on plastic or carbon-coated nickel grid. Float on a drop of water for 5 - 10 minutes.
2. Incubate with 1 % solution of bovine serum albumin in PBS buffer at pH 7.4 for 5 minutes to block non-specific protein binding sites.
3. Rinse with PBS-BSA (1 min).
4. Incubate with UNDECAGOLD conjugate diluted 1/40 - 1/200 in PBS-BSA with 1 % normal serum from the same species as the UNDECAGOLD labeled antibody, for 10 minutes to 1 hour at room temperature.
5. Rinse with PBS (3 X 1 min).
6. Postfix with 1 % glutaraldehyde in PBS at room temperature (3 mins).

7. Rinse in deionized water for (2 X 5 min).
8. If desired, contrast sections with uranyl acetate and/or lead citrate before examination.

Silver enhancement may also be used to render the UNDECAGOLD particles more easily visible (see below), especially if stains such as uranyl acetate or lead citrate are applied. If used, it should be completed before these stains are applied.

PBS-BSA Buffer:

20 mM phosphate
150 mM NaCl
pH 7.4
0.5% BSA
0.1% gelatin (high purity)

Optional, may reduce background:

0.5 M NaCl
0.05% Tween 20

PBS Buffer:

20 mM phosphate
150 mM NaCl
pH 7.4

SPECIAL CONSIDERATIONS FOR VIEWING UNDECAGOLD IN THE ELECTRON MICROSCOPE

UNDECAGOLD is the smallest gold probe commercially available, being just 0.8 nm in diameter. A high resolution instrument such as a Scanning Transmission Electron Microscope (STEM) is required for visualization; in a conventional TEM the UNDECAGOLD particles are not visible. With careful work, however, UNDECAGOLD may be seen directly in the STEM. However, achieving the high resolution necessary for this work may require new demands on your equipment and technique. Several suggestions follow:

1. Before you start a project with UNDECAGOLD it is helpful to see it so you know what to look for. Dilute the UNDECAGOLD stock 1:5 in methanol and apply 4 μ l to a grid for 1 minute. Allow to dry.
2. View UNDECAGOLD using a full width scan of 128 nm or less; this will give sufficient magnification for visualization.
3. UNDECAGOLD is sensitive to beam damage (contrary to NANOGOLD[®] which is very beam-resistant); the behavior of UNDECAGOLD in the STEM has been described in the literature.² Image at approximately 200 e⁻ Å⁻².
4. In order to operate at high magnification, thin carbon film over fenestrated holey film is recommended. Many plastic supports are unstable under these conditions of high magnification/high beam current and carbon is therefore preferred. Contrast is best using thinner films.

SILVER ENHANCEMENT OF UNDECAGOLD FOR EM

UNDECAGOLD will nucleate silver deposition resulting in a dense particle 2-20 nm in size or larger depending on development time. However, silver enhancement will be slower and much less uniform than with larger gold particles such as NANOGOLD[®]. If specimens are to be embedded, silver enhancement is usually performed after embedding, although it may be done first. It must be completed before any staining reagents such as osmium tetroxide, lead citrate or uranyl acetate are applied, since these will nucleate silver deposition in the same manner as gold and produce non-specific staining.

Our LI SILVER silver enhancement system is convenient and not light sensitive, and suitable for all applications. Improved results in the EM may be obtained using HQ SILVER, which is formulated to give slower, more controllable particle growth and uniform particle size distribution.

Specimens must be thoroughly rinsed with deionized water before silver enhancement reagents are applied. This is because the buffers used for antibody incubations and washes contain chloride ions and other anions which form insoluble precipitates with silver. These are often light-sensitive and will give non-specific staining. To prepare the developer, mix equal amounts of the enhancer and initiator immediately before use. UNDECAGOLD will nucleate silver deposition resulting in a dense particle 2-20 nm in size or larger depending on development time. Use nickel grids (not copper).

The procedure for immunolabeling should be followed up to step 6 as described above. Silver enhancement is then performed as follows:

1. Rinse with deionized water (2 X 5 mins).
2. Float grid with specimen on freshly mixed developer for 1-4 minutes, or as directed in the instructions for the silver reagent. More or less time can be used to control particle size. A series of different development times should be tried, to find the optimum time for your experiment.
3. Rinse with deionized water (3 X 1 min).
4. Mount as usual.

LIGHT MICROSCOPY WITH UNDECAGOLD

For most light microscopy applications we recommend NANOGOLD[®] conjugates, since these usually give more intense, specific signals. UNDECAGOLD must be developed with a silver enhancement reagent before it is visible in the light microscope. Our LI SILVER is convenient for this as it is not light sensitive and the degree of enhancement may be monitored readily.

Samples must be rinsed with deionized water before silver enhancement. This is because the reagent contains silver ions in solution, which react to form a precipitate with chloride, phosphate and other anions which are components of buffer solutions. The procedure for immunolabeling with UNDECAGOLD and silver enhancement is given below.

1. Spin cells onto slides using Cytospin, or use paraffin section.
2. Incubate with 1 % solution of bovine serum albumin in PBS (PBS-BSA) for 10 minutes to block non-specific protein binding sites.
3. Incubate with primary antibody, diluted at usual working concentration in PBS-BSA (1 hour or usual time)
4. Rinse with PBS-BSA (3 X 2 min).
5. Incubate with UNDECAGOLD reagent diluted 1/40 - 1/200 in PBS-BSA with 1 % normal serum from the same species as the UNDECAGOLD reagent, for 1 hour at room temperature.
6. Rinse with PBS (3 X 5 min).
7. Postfix with 1 % glutaraldehyde in PBS at room temperature (3 mins).
8. Rinse with deionized water (3 X 1 min).
9. Develop specimen with freshly mixed developer for 5-20 minutes, or as directed in the instructions for the silver reagent. More or less time can be used to control intensity of signal. A series of different development times may be used, to find the optimum enhancement for your experiment; generally a shorter antibody incubation time will require a longer silver development time.
10. Rinse with deionized water (2 X 5 mins).
11. The specimen may now be stained if desired before examination, with usual reagents.

PBS-BSA Buffer:

20 mM phosphate
150 mM NaCl
pH 7.4
0.5% BSA
0.1% gelatin (high purity)

Optional, may reduce background:

0.5 M NaCl
0.05% Tween 20

PBS Buffer:

20 mM phosphate
150 mM NaCl
pH 7.4

To obtain an especially dark silver signal, the silver enhancement may be repeated with a freshly mixed portion of developer.

IMMUNOBLOTTING

For immunoblotting we recommend our NANOGOLD[®] conjugates, since these give more sensitive detection than UNDECAGOLD conjugates with a higher signal-to-noise ratio upon silver enhancement. The basic procedure for gold immunoblotting has been

described by Moeremans et al⁵, which may be followed. For best results, the membrane should be hydrated before use by simmering in gently boiling water for 15 minutes. Best results are obtained when the antigen is applied using a 1 µl capillary tube. A suitable procedure for immunoblots is as follows:

1. Spot 1 µl dilutions of the antigen in buffer 4 onto hydrated nitrocellulose membrane. Use an antigen concentration range from 100 to 0.01 pg / µl.
2. Block with buffer 1 for 30 minutes at 37°C.
3. Incubate with primary antibody according to usual procedure (usually 1 or 2 hours).
4. Rinse with buffer 1 (3 X 10 mins).
5. Incubate with a 1/100 to 1/200 dilution of the UNDECAGOLD reagent in buffer 2 for 2 hours at room temperature.
6. Rinse with buffer 3 (3 X 5 mins), then buffer 4 (2 X 5 mins).
7. OPTIONAL (may improve sensitivity): Postfix with glutaraldehyde, 1 % in buffer 4 (10 mins).
8. Rinse with deionized water (2 X 5 mins).
9. OPTIONAL (may reduce background): Rinse with 0.05 M EDTA at pH 4.5 (5 mins).
10. Develop with freshly mixed silver developer for 20-25 minutes or as directed in the instructions for the silver reagent, twice. Rinse thoroughly with deionized water between developments to remove all the reagent.
11. Rinse several times with deionized water.

CAUTION: UNDECAGOLD particles degrade upon exposure to concentrated thiols such as β-mercaptoethanol or dithiothreitol. If such reagents must be used, concentrations should be kept below 1 mM and exposure restricted to 10 minutes or less.

Buffer 1: 20 mM phosphate
150 mM NaCl
pH 7.4
4% BSA (bovine serum albumin)
2 mM sodium azide (NaN₃)

Buffer 3: 20 mM phosphate
150 mM NaCl
pH 7.4
0.8% BSA (bovine serum albumin)
2 mM sodium azide (NaN₃)

Buffer 2: 20 mM phosphate
150 mM NaCl
pH 7.4
0.8% BSA
1% normal serum; use serum of the host animal
for the UNDECAGOLD-conjugated
antibody
0.1% gelatin (Type B, approx. 60 bloom)
Optional, may reduce background:
0.5 M NaCl
0.05% Tween 20

Buffer 4 (PBS):
20 mM phosphate
150 mM NaCl
pH 7.4

Other procedures may be used; for example the UNDECAGOLD reagent may be used as a tertiary labeled antibody, or a custom UNDECAGOLD conjugate may be the primary antibody. If additional antibody incubation steps are used, rinse with buffer 3 (3 X 10 mins) after incubation.

REFERENCES

1. Hainfeld, J. F.; in "Colloidal Gold: Principles, Methods and Applications;" M. A. Hayat, ed.; Vol. 2, p. 413; Academic Press, San Diego, CA (1991).
2. Parham, P, *J. Immunology*, 131, 2895 (1983).
3. Grasseti, D. R., and Murray, J. F. Jr.; *Arch. Biochem. Biophys.*, 119, 41 (1967).
4. J. E. Beesley, in "Colloidal Gold: Principles, Methods and Applications," M. A. Hayat, ed., Academic Press, New York, 1989; Vol. 1, pp421-425.
5. Moeremans, M. et al., *J. Immunol. Meth.* 74, 353 (1984).

Technical Assistance Available.

For a complete list of references citing this product, please visit our world-wide-web site at <http://www.nanoprobes.com/Ref.html>.